
WO 2005/045035 PCT/US2004/026930

1/25

## Figure 1



= SOLID SUPPORT

R = TERMINAL PROTECTING GROUP FOR EXAMPLE: DIMETHOXYTRITYL (DMT)


= CLEAVABLE LINKER

(FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR

(NVERTED DEOXYABASIC SUCCINATE)

= CLEAVABLE LINKER


(FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR INVERTED DEOXYABASIC SUCCINATE)




INVERTED DEOXYABASIC SUCCINATE LINKAGE

GLYCERYL SUCCINATE LINKAGE

Figure 2





### 4/25

## Figure 4

		·	
A		SENSE STRAND (SEQ ID NO 303) ALL POSITIONS RIBONUCLEOTIDE EXCEPT POSITIONS (N N)	
	5'-	B-NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	-3'
	<b>3'-</b>	L-(N <sub>S</sub> N) NNNNNNNNNNNNNNNNNN	-5'
		ANTISENSE STRAND (SEQ ID NO 304) ALL POSITIONS RIBONUCLEOTIDE EXCEPT POSITIONS (N N)	J
	ALL	SENSE STRAND (SEQ ID NO 305) PYRIMIDINES = 2'-FLUORO AND ALL PURINES = 2'-OME EXCEPT POSITIONS (	(и и
	5'-	NNNNNNNNNNNNNNNN(N <sub>S</sub> N)	-3'
B	₹ 3'-	L-(N <sub>S</sub> N) NNNNNNNNNNNNNNNNNNN	-5'
	ALL	ANTISENSE STRAND (SEQ ID NO 306)  PYRIMIDINES = 2'-FLUORO AND ALL PURINES = 2'-OME EXCEPT POSITIONS (N	(N I
	Ċ		<u> </u>
		SENSE STRAND (SEQ ID NO 307) ALL PYRIMIDINES = 2'-O-ME OR 2'-FLUORO EXCEPT POSITIONS (N N)	
	5'-	B-N N N N N N N N N N N N N N N N N N (N N)-B	-3'
C	3'-	L-(N <sub>S</sub> N) N N N N N N N N N N N N N N N N N N	-5' [
		ANTISENSE STRAND (SEQ ID NO 308) ALL PYRIMIDINES = 2'-FLUORO EXCEPT POSITIONS (N N)	J
D	(ALL B	SENSE STRAND (SEQ ID NO 309) YRIMIDINES = 2'-FLUORO EXCEPT POSITIONS (N N) AND ALL PURINES = 2'-DE	(yxo
	5'-	B-NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	-3'
	₹3'-	L-(N <sub>S</sub> N) NNNNNNNNNNNNNNNNN	-5'
		ANTISENSE STRAND (SEQ ID NO 306) PYRIMIDINES = 2'-FLUORO AND ALL PURINES = 2'-O-ME EXCEPT POSITIONS (	NN)
		SENSE STRAND (SEQ ID NO 310)	Š
		SENSE STRAND (SEQ ID NO 310) ALL PYRIMIDINES = 2'-FLUORO EXCEPT POSITIONS (N N)	
E	5'-	B-NNNNNNNNNNNNNNNNNNNNNNNNNN-B	(
	3'-	L-(N <sub>S</sub> N) NNNNNNNNNNNNNNNNN	-5'
	ALL	ANTISENSE STRAND (SEQ ID NO 306) PYRIMIDINES = 2'-FLUORO AND ALL PURINES = 2'-O-ME EXCEPT POSITIONS (I	(N N
		SENSE STRAND (SEQ ID NO 309)	)
	1	YRIMIDINES = $2'$ -FLUORO EXCEPT POSITIONS (N N) AND ALL PURINES = $2'$ -DE	
F	<b>∫</b> 5'-	B-NNNNNNNNNNNNNNNNN(NN)-B	-3'
	3'-	L-(N <sub>S</sub> N) NNNNNNNNNNNNNNNNN	-5'
	ALLP	ANTISENSE STRAND (SEQ ID NO 311) 'YRIMIDINES = 2'-FLUORO EXCEPT POSITIONS (N N) AND ALL PURINES = 2'-DE	OXY

POSITIONS (NN) CAN COMPRISE ANY NUCLEOTIDE, SUCH AS DEOXYNUCLEOTIDES (eg. THYMIDINE) OR UNIVERSAL BASES

B = ABASIC, INVERTED ABASIC, INVERTED NUCLEOTIDE OR OTHER TERMINAL CAP THAT IS OPTIONALLY PRESENT

L = GLYCERYL MOIETY THAT IS OPTIONALLY PRESENT

S = PHOSPHOROTHIOATE OR PHOSPHORODITHIOATE THAT IS OPTIONALLY PRESENT

5/25

## Figure 5


		SENSE STRAND (SEQ ID NO 312)	
A	5'- 3'-	B-GGCUGCUG GCAUGGGUGCU <i>TT</i> -B L-T <sub>S</sub> TCCGACGACCGUACCCACGA ANTISENSE STRAND (SEQ ID NO 313)	-3' -5'
В	5'- 3'-	SENSE STRAND (SEQ ID NO 314)  ggcugcuggguggugcuT <sub>S</sub> T  L-T <sub>S</sub> T c c g a c g a c c g u a c c c a c g a  ANTISENSE STRAND (SEQ ID NO 315)	-3' -5'
C	5'- 3'-	SENSE STRAND (SEQ ID NO 316)  B-GGcuGcuGcuGGCuGGuGcuTT-B  L-T <sub>S</sub> TccGAAGAACCGUACCAACAA  ANTISENSE STRAND (SEQ ID NO 317)	-3' -5'
D	5'- 3'-	SENSE STRAND (SEQ ID NO 318)  B-G G c u G c u G G c A u G G G u G c u T T-B  L-T <sub>S</sub> T c c g a c g a c c g u a c c c a c g a  ANTISENSE STRAND (SEQ ID NO 315)	-3' -5'
E	5'- 3'-	SENSE STRAND (SEQ ID NO 319)  B-G G c u G c u G G c A u G G G u G c u T T-B  L-T <sub>S</sub> T c c g <u>a</u> c g <u>a</u> c c g u <u>a</u> c c c <u>a</u> c g <u>a</u> ANTISENSE STRAND (SEQ ID NO 315)	-3' -5'
$\mathbf{F}$	5'- 3'-	SENSE STRAND (SEQ ID NO 318)  B-GGcuGcuGGCAuGGGuGcuTT-B  L-T <sub>S</sub> TccGAcGAcGACGA  ANTISENSE STRAND (SEQ ID NO 320)	-3' -5'


lower case = 2'-O-Methyl or 2'-deoxy-2'-fluoro italic lower case = 2'-deoxy-2'-fluoro


<u>underline</u> = 2'-O-methyl ITALIC UPPER CASE = DEOXY B = ABASIC, INVERTED ABASIC, INVERTED NUCLEOTIDE OR OTHER TERMINAL CAP THAT IS OPTIONALLY PRESENT

S = PHOSPHOROTHIOATE ORPHOSPHORODITHIOATE OPTIONALLY PRESENT


L = GLYCERYL MOIETY, OR B, OPTIONALLY PRESENT










ANTISENSE


















n = 0, 1, 2, 3, 4








Identify efficacious target sites based on siRNA sequence Sequence siRNA Clone oligos into vector Figure 9: Target site Selection using siRNA Select cells exhibiting desired phenotype Ш nence Synthesize oligos encoding siRNA against Target RNA seq Transduce target cells




R = O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, or aralkyl B = Independently any nucleotide base, either naturally occurring or chemically modified, or optionally H (abasic).

Figure 11: Modification Strategy




# Figure 12: Phosphorylated siNA constructs



modifications herein


## sphate modifications




oligonucleotide constructs that utilize or repeat sequences Palindrome Figure 14A: Duplex forming

Sequence of (ii) to 3'-end of complementary Target Nucleic Acid sequence of (i) above Design Complementary Sequence to the containing palindrome/repeat sequence Self assembly of self complementary Identify Target Nucleic Acid sequence (e.g., 14 to 24 nucleotides in length) strands to form duplex construct Non-palindromic Complementary Append inverse sequence of the at 5'-end (dashed portion) sednence 111 Ñ Ŝ က် ကိ က်  $(\tilde{\mathbb{N}})$  $\equiv$ 


Figure 14B: Example of a duplex forming oligonucleotide sequence




forming oligonucleotide sequence epeat sequence, self assembly that utilizes a palindrome or re Figure 14C: Example of a duplex



forming oligonucleotide sequence that utilizes a palindrome or repeat sequence, self assembly and inhibition Figure 14D: Example of a duplex



oligonucleotide constructs that utilize rome or repeat sequences artificial palind Figure 15: Duplex forming




Identify Target Nucleic Acid sequence (e.g., 14 to 24 nucleotides in length)

Design Complementary Sequence and utilize modified nucleotides (shown as X, Y) that interact with a portion of the target sequence and result in the formation of a palindrome/repeat sequence (e.g., 2 to 12 nucleotides) at 3'-end (dashed portion)

Append inverse sequence of Complementary region to 3'-end of palindrome/repeat sequence

က်



Hybridize self complementary strands to form duplex siNA construct






Figure 17: Examples of hairpin multifunctional siNA constructs Complementary Region 2 Complementary Region 1 with distinct complementary regions Complementary Region Complementary Region က် က် က်က် **m** 

distinct complementary regions and a self complementary/palindrome region Inded multifunctional siNA constructs with Figure 18: Examples of double stra



multifunctional siNA constructs with Figure 19: Examples of hairpin







Figure 21: Example of multifunctional siNA targeting two regions within the same target nucleic acid sequence Region 2 RISC Processing Region 1 X = cleavage ත ත





## This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:				
☐ BLACK BORDERS				
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES				
FADED TEXT OR DRAWING				
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING				
☐ SKEWED/SLANTED IMAGES				
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS				
☐ GRAY SCALE DOCUMENTS				
LINES OR MARKS ON ORIGINAL DOCUMENT				
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY				
OTHER:				

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.